

# CS228 Logic for Computer Science 2021

## Lecture 15: Handling first-order logic

Instructor: Ashutosh Gupta

IITB, India

Compile date: 2021-02-10

# Topic 15.1

## Supporting definitions

## Clubbing similar quantifiers

If we have a chain of **same quantifier** then we write the quantifier **once** followed by the list of variables.

### Example 15.1

- ▶  $\forall z, x. \exists y. G(x, y, z) = \forall z. (\forall x. (\exists y. G(x, y, z)))$
- ▶  $\exists z, x, y. G(x, y, z) = \exists z. (\exists x. (\exists y. G(x, y, z)))$

# Subterm and subformulas

## Definition 15.1

A term  $t$  is *subterm* of term  $t'$ , if  $t$  is a substring of  $t'$ .

## Exercise 15.1

- ▶ Is  $f(x)$  a subterm of  $g(f(x), y)$ ?
- ▶ Is  $c$  a subterm of  $c$ ?
- ▶  $x$  is a subterm of  $f(x)$

## Definition 15.2

A formula  $F$  is *subformula* of formula  $F'$ , if  $F$  is a substring of  $F'$ .

## Example 15.2

- ▶  $G(x, y, z)$  is a subformula of  $\forall z, x. \exists y. G(x, y, z)$
- ▶  $P(c)$  is a subformula of  $P(c)$
- ▶  $\exists y. G(x, y, z)$  is a subformula of  $\forall z, x. \exists y. G(x, y, z)$

# Closed terms and quantifier free

## Definition 15.3

A *closed term* is a term without any variable. Let  $\hat{T}_{\mathbf{S}}$  be the set of closed  $\mathbf{S}$ -terms.

Sometimes closed terms are also referred as *ground terms*.

## Example 15.3

Let  $\mathbf{F} = \{f/1, c/0\}$ .  $f(c)$  is a closed term, and  $f(x)$  is not, where  $x$  is a variable.

## Exercise 15.2

Which of the following terms are closed with respect to  $\mathbf{F} = \{f/1, g/2, c/0\}$ ?

▶  $g(c, y)$

▶  $c$

▶  $x$

▶  $f(g(c, c))$

# Quantifier-free

## Definition 15.4

A formula  $F$  is *quantifier-free* if there are no quantifiers in  $F$ .

## Example 15.4

$H(c)$  is a quantifier-free formula and  $\forall x.H(x)$  is not a quantifier-free formula.

## Exercise 15.3

For signature  $(\{f/1, c/0\}, \{H/1\})$ , which of the following are quantifier-free?

- ▶  $\forall x.H(y)$
- ▶  $f(c)$
- ▶  $H(y) \vee \perp$
- ▶  $H(f(c))$

# Free variables

## Definition 15.5

A variable  $x \in \text{Vars}$  is *free* in formula  $F$  if

- ▶  $F \in A_S$ :  $x$  occurs in  $F$ ,
- ▶  $F = \neg G$ :  $x$  is free in  $G$ ,
- ▶  $F = G \circ H$ :  $x$  is free in  $G$  or  $H$ , for some binary operator  $\circ$ , and
- ▶  $F = \exists y.G$  or  $F = \forall y.G$ :  $x$  is free in  $G$  and  $x \neq y$ .

Let  $FV(F)$  denote the set of free variables in  $F$ .

## Exercise 15.4

Is  $x$  free?

- |          |                                      |
|----------|--------------------------------------|
| ▶ $H(x)$ | ▶ $\forall x.H(x)$                   |
| ▶ $H(y)$ | ▶ $x = y \Rightarrow \exists x.G(x)$ |

# Sentence

## Definition 15.6

In  $\forall x.(G)$ , we say the quantifier  $\forall x$  has *scope*  $G$  and *bounds*  $x$ .

In  $\exists x.(G)$ , we say the quantifier  $\exists x$  has *scope*  $G$  and *bounds*  $x$ .

## Definition 15.7

A formula  $F$  is a *sentence* if it has no free variable.

## Exercise 15.5

Which of the following formulas are sentence(s)?

▶  $H(x)$

▶  $\forall x.H(x)$

▶  $x = y \Rightarrow \exists x.G(x)$

▶  $\forall x.\exists y. x = y \Rightarrow \exists x.G(x)$



## Topic 15.2

### Understanding FOL semantics

## No free variables

### Definition 15.8

Let  $t$  be a closed term.  $m(t) \triangleq m^\nu(t)$  for any  $\nu$ .

If  $F$  is a sentence,  $\nu$  has no influence in the satisfaction relation.(why?)

For sentence  $F$ , we say

- ▶  $F$  is *true* in  $m$  if  $m \models F$
- ▶ Otherwise,  $F$  is *false* in  $m$ .

## Why nonempty domain?

We are required to have **nonempty domain** in the model. Why?

### Example 15.5

Consider formula  $\forall x.(H(x) \wedge \neg H(x))$ .

Should any model satisfy the formula?

*Noooooooooo..*

But, if we allow  $m = \{\emptyset; H_m = \emptyset\}$  then


$$m \models \forall x.(H(x) \wedge \neg H(x)).$$

Due to this counter-intuitive behavior, the **empty domain** is disallowed.

## Example: graph models

### Example 15.6

Consider  $\mathbf{S} = (\{\}, \{E/2\})$  and  $m = (\{a, b\}; \{(a, a), (a, b)\})$ .  
 $m$  may be viewed as the following graph.



$$m, \{x \rightarrow a\} \models E(x, x) \wedge \exists y. (E(x, y) \wedge \neg E(y, y))$$

### Exercise 15.6

Give another model and assignment that satisfies the above formula

## Example : counting

### Example 15.7

Consider  $\mathbf{S} = (\{\}, \{E/2\})$

*The following sentence is false in all the models with one element domain*

$$\forall x. \neg E(x, x) \wedge \exists x \exists y. E(x, y)$$

## Exercise: counting

### Exercise 15.7

*Give a sentence that is true only in the models with more than two elements*

### Exercise 15.8

- a. Give a sentence that is true only in infinite models*
- b. Do only finite models satisfy the negation of the sentence in (a)? If not, give an example of infinite model.*

### Exercise 15.9

- a. Give a sentence that is true only in models with less than or equal to two element domains.*
- b. Can you answer (a) without using  $=$ ?*

## A limit: Impossibility of expressing finite

### Theorem 15.1

*No FOL sentence can express that all satisfying models are finite.*

Commentary: Proof of the above is not part of this course.

## Topic 15.3

### Substitution



# Substitution

In first-order logic, we have terms and formulas. We need a more elaborate notion of substitution for terms.

## Definition 15.9

A *substitution*  $\sigma$  is a map from  $\text{Vars} \rightarrow T_S$ . We will write  $t\sigma$  to denote  $\sigma(t)$ .

## Definition 15.10

We say  $\sigma$  has *finite support* if only finite variables do not map to themselves.  $\sigma$  with *finite support* is denoted by  $[t_1/x_1, \dots, t_n/x_n]$  or  $\{x_1 \mapsto t_1, \dots, x_n \mapsto t_n\}$ .

We may write a formula as  $F(x_1, \dots, x_k)$ , where variables  $x_1, \dots, x_k$  play a special role in  $F$ .

Let  $F(t_1, \dots, t_n)$  be  $F[t_1/x_1, \dots, t_n/x_n]$ .

**Commentary:** We have seen substitution in propositional logic. Now in FOL, substitution is not a simple matter. We were replacing a formula by another. Now we need another kind of substitution that replaces terms.

# Substitution on terms

## Definition 15.11

For  $t \in T_{\mathbf{S}}$ , let the following naturally define  $t\sigma$  as extension of  $\sigma$ .

- ▶  $c\sigma \triangleq c$
- ▶  $(f(t_1, \dots, t_n))\sigma \triangleq f(t_1\sigma, \dots, t_n\sigma)$

## Example 15.8

Consider  $\sigma = \{x \mapsto f(x, y), y \mapsto f(y, x)\}$

- ▶  $x\sigma = f(x, y)$
- ▶  $f(x, y)\sigma = f(f(x, y), f(y, x))$
- ▶  $(f(x, y)\sigma)\sigma = ?$

# Substitution on atoms

We further extend the substitution  $\sigma$  to atoms.

## Definition 15.12

For  $F \in A_{\mathbf{S}}$ ,  $F\sigma$  is defined as follows.

- ▶  $\top\sigma \triangleq \top$
- ▶  $\perp\sigma \triangleq \perp$
- ▶  $P(t_1, \dots, t_n)\sigma \triangleq P(t_1\sigma, \dots, t_n\sigma)$
- ▶  $(t_1 = t_2)\sigma \triangleq t_1\sigma = t_2\sigma$

# Substitution projection

Sometimes, we may need to remove variable  $x$  from the support of  $\sigma$ .

## Definition 15.13

Let  $\sigma_x = \sigma[x \mapsto x]$ .

## Example 15.9

Consider  $\sigma = \{x \mapsto f(x, y), y \mapsto f(y, x)\}$ .  $\sigma_x = \{y \mapsto f(y, x)\}$

Commentary: The need of the definition will be clear soon.

## Substitution in formulas (Incorrect)

Now we extend the substitution  $\sigma$  to all the formulas.

### Definition 15.14

For  $F \in P_S$ ,  $F\sigma$  is defined as follows.

- ▶  $(\neg G)\sigma \triangleq \neg(G\sigma)$
- ▶  $(G \circ H)\sigma \triangleq (G\sigma) \circ (H\sigma)$  for some binary operator  $\circ$
- ▶  $(\forall x.G)\sigma \triangleq \forall x.(G\sigma_x)$
- ▶  $(\exists x.G)\sigma \triangleq \exists x.(G\sigma_x)$

### Example 15.10

- ▶  $(P(x) \Rightarrow \forall x.Q(x))\{x \mapsto y\} = (P(y) \Rightarrow \forall x.Q(x))$
- ▶  $(\exists y. x \neq y)\{x \mapsto z\} = (\exists y. z \neq y)$
- ▶  $(\exists y. x \neq y)\{x \mapsto y\} = (\exists y. y \neq y)$  ☹️ Undesirable!!!

Some substitutions should be disallowed.

**Commentary:** The above naïve definition of the substitution in formulas is incorrect. In the next slide, we present the correct definition.

## Substitution in formulas(Correct)

### Definition 15.15

$\sigma$  is *suitable* with respect to formula  $G$  and variable  $x$  if for all  $y \neq x$ , if  $y \in FV(G)$  then  $x$  does not occur in  $y\sigma$ .

Now we *correctly* extend the substitution  $\sigma$  to all formulas.

### Definition 15.16

For  $F \in P_S$ ,  $F\sigma$  is defined as follows.

- ▶  $(\neg G)\sigma \triangleq \neg(G\sigma)$
- ▶  $(G \circ H)\sigma \triangleq (G\sigma) \circ (H\sigma)$  for some binary operator  $\circ$
- ▶  $(\forall x.G)\sigma \triangleq \forall x.(G\sigma_x)$ , where  $\sigma$  is suitable with respect to  $G$  and  $x$
- ▶  $(\exists x.G)\sigma \triangleq \exists x.(G\sigma_x)$ , where  $\sigma$  is suitable with respect to  $G$  and  $x$

It is not a true restriction.  
We will see later.

# Composition

## Definition 15.17

Let  $\sigma_1$  and  $\sigma_2$  be substitutions. The **composition**  $\sigma_1\sigma_2$  of the substitutions is defined as follows.

$$\text{For each } x \in \text{Vars}, x(\sigma_1\sigma_2) \triangleq (x\sigma_1)\sigma_2.$$

## Example 15.11

- ▶  $\sigma_1 = \{x \mapsto f(x, y)\}$  and  $\sigma_2 = \{y \mapsto c\}$ .  $\sigma_1\sigma_2 = \{x \mapsto f(x, c), y \mapsto c\}$ .
- ▶  $\sigma_1 = \{x \mapsto y\}$  and  $\sigma_2 = \{y \mapsto x\}$ .  $\sigma_1\sigma_2 = \{x \mapsto x, y \mapsto x\} = \{y \mapsto x\}$ .

## Exercise 15.10

Show  $\sigma_1(\sigma_2\sigma_3) = (\sigma_1\sigma_2)\sigma_3$ , i.e., substitution is associative.

**Commentary:** Type check composition definition. Convince yourself that composition is well-defined. Solution for exercise: Consider variable  $x$ .  $(x\sigma_1)(\sigma_2\sigma_3) = ((x\sigma_1)\sigma_2)\sigma_3 = (x(\sigma_1\sigma_2))\sigma_3$

# Composition works on terms and atoms

## Theorem 15.2

For each  $t \in T_{\mathbf{S}}$ ,  $t(\sigma_1\sigma_2) = (t\sigma_1)\sigma_2$

Proof.

Proved by trivial structural induction. □

## Theorem 15.3

For each  $F \in A_{\mathbf{S}}$ ,  $F(\sigma_1\sigma_2) = (F\sigma_1)\sigma_2$

Proof.

Proved by trivial structural induction. □



# Substitution composition on formulas

## Theorem 15.4

if  $F\sigma_1$  and  $(F\sigma_1)\sigma_2$  are defined then  $(F\sigma_1)\sigma_2 = F(\sigma_1\sigma_2)$

**Proof.**

We prove it by induction. Non-quantifier cases are simple structural induction.

Assume  $F = \forall x.G$

Since  $F\sigma_1$  is defined,  $G\sigma_{1x}$  is defined. Since  $(F\sigma_1)\sigma_2$  is defined,  $(G\sigma_{1x})\sigma_{2x}$  is defined (why?).

By induction hypothesis,  $(G\sigma_{1x})\sigma_{2x} = G(\sigma_{1x}\sigma_{2x})$

**claim:**  $G(\sigma_{1x}\sigma_{2x}) = G(\sigma_1\sigma_2)_x$

Choose  $y \in FV(G)$  and  $y \neq x$

$$y(\sigma_{1x}\sigma_{2x}) = \underbrace{((y\sigma_{1x})\sigma_{2x})}_{\text{Def. substitution}} = \underbrace{((y\sigma_1)\sigma_{2x})}_{y \neq x} = \underbrace{((y\sigma_1)\sigma_2)}_{x \notin FV(y\sigma_1) \text{ (why?)}} = y(\sigma_1\sigma_2) = y(\sigma_1\sigma_2)_x$$

$$(\forall x.G\sigma_1)\sigma_2 = (\forall x.G(\sigma_{1x}\sigma_{2x})) = (\forall x.G(\sigma_1\sigma_2)_x) = F(\sigma_1\sigma_2)$$

□

## Topic 15.4

### Problems

# Properties of FOL

## Exercise 15.11

If  $x, y \notin \text{Vars}(F(z))$ , then  $\forall x.F(x) \Leftrightarrow \forall y.F(y)$

## Exercise 15.12

Let us suppose  $x$  does not occur in formula  $G$ . Show that the following formulas are valid.

- ▶  $\exists x.G \Leftrightarrow G$
- ▶  $\forall x.G \Leftrightarrow G$
- ▶  $(\forall x.F(x) \vee G) \Leftrightarrow \forall x.(F(x) \vee G)$
- ▶  $(\forall x.F(x) \wedge G) \Leftrightarrow \forall x.(F(x) \wedge G)$
- ▶  $(\exists x.F(x) \vee G) \Leftrightarrow \exists x.(F(x) \vee G)$
- ▶  $(\exists x.F(x) \wedge G) \Leftrightarrow \exists x.(F(x) \wedge G)$

## Encode mod $k$

### Exercise 15.13

*Give an FOL sentence that encodes that there are  $n$  elements in any satisfying model, such that  $n \bmod k = 0$  for a given  $k$ .*

# Unique quantifier

## Exercise 15.14

*We could consider enriching the language by the addition of a new quantifier. The formula  $\exists! x.F$  (read “there exists a unique  $x$  such that  $F$ ”) is to be satisfied in model  $m$  and assignment  $\nu$  iff there is one and only one  $d \in D_m$  such that  $m, \nu[x \rightarrow d] \models F$ . Show that this apparent enrichment does not increase expressive power of FOL.*

## Exercise: equality propagation

### Exercise 15.15

*Which of the following equivalences are correct?*

- ▶  $\exists x, x'. (x' = x \wedge F(x, x')) \equiv \exists x. F(x, x)$
- ▶  $\exists x, x'. (x' = x \Rightarrow F(x, x')) \equiv \exists x. F(x, x)$
- ▶  $\forall x, x'. (x' = x \wedge F(x, x')) \equiv \forall x. F(x, x)$
- ▶  $\forall x, x'. (x' = x \Rightarrow F(x, x')) \equiv \forall x. F(x, x)$

## Topic 15.5

Extra slides: not-so-useful definitions

# Bounded variables

## Definition 15.18

A variable  $x \in \text{Vars}$  is *bounded* in formula  $F$  if

- ▶  $F = \neg G$ :  $x$  is bounded in  $G$ ,
- ▶  $F = G \circ H$ :  $x$  is bounded in  $G$  or  $H$ , for some binary operator  $\circ$ , and
- ▶  $F = \exists y.G$  or  $F = \forall y.G$ :  $x$  is bounded in  $G$  or  $x$  is equal to  $y$ .

Let  $\text{bnd}(F)$  denote the set of bounded variables in  $F$ .

## Exercise 15.16

Is  $x$  bounded?

- |          |                                      |
|----------|--------------------------------------|
| ▶ $H(x)$ | ▶ $\forall x.H(x)$                   |
| ▶ $H(y)$ | ▶ $x = y \Rightarrow \exists x.G(x)$ |



End of Lecture 15