

CS228 Logic for Computer Science 2022

Lecture 15: FOL - formal proofs

Instructor: Ashutosh Gupta

IITB, India

Compile date: 2022-02-26

Topic 15.1

Formal proofs

Consequence to derivation

We also need the formal proof system for FOL.

Let us suppose for a (in)finite set of formulas Σ and a formula F , we have $\Sigma \models F$.

Similar to propositional logic, we will now again develop a system of “derivations”. We derive the following **statements**.

$$\Sigma \vdash F$$

Formal rules for FOL

- ▶ The old rules will continue to work
- ▶ We need new rules for.....
- ▶ Let us see how do we develop those!

quantifiers and equality

Rules for propositional logic stays!

$$\text{ASSUMPTION} \frac{}{\Sigma \vdash F} F \in \Sigma \quad \text{MONOTONIC} \frac{\Sigma \vdash F}{\Sigma' \vdash F} \Sigma \subseteq \Sigma' \quad \text{DOUBLENEG} \frac{\Sigma \vdash F}{\Sigma \vdash \neg\neg F}$$

$$\wedge\text{-INTRO} \frac{\Sigma \vdash F \quad \Sigma \vdash G}{\Sigma \vdash F \wedge G} \quad \wedge\text{-ELIM} \frac{\Sigma \vdash F \wedge G}{\Sigma \vdash F} \quad \wedge\text{-SYMM} \frac{\Sigma \vdash F \wedge G}{\Sigma \vdash G \wedge F}$$

$$\vee\text{-INTRO} \frac{\Sigma \vdash F}{\Sigma \vdash F \vee G} \quad \vee\text{-SYMM} \frac{\Sigma \vdash F \vee G}{\Sigma \vdash G \vee F} \quad \vee\text{-DEF} \frac{\Sigma \vdash F \vee G}{\Sigma \vdash \neg(\neg F \wedge \neg G)}^*$$

$$\vee\text{-ELIM} \frac{\Sigma \vdash F \vee G \quad \Sigma \cup \{F\} \vdash H \quad \Sigma \cup \{G\} \vdash H}{\Sigma \vdash H}$$

$$\Rightarrow\text{-INTRO} \frac{\Sigma \cup \{F\} \vdash G}{\Sigma \vdash F \Rightarrow G} \quad \Rightarrow\text{-ELIM} \frac{\Sigma \vdash F \Rightarrow G \quad \Sigma \vdash F}{\Sigma \vdash G} \quad \Rightarrow\text{-DEF} \frac{\Sigma \vdash F \Rightarrow G}{\Sigma \vdash \neg F \vee G}^*$$

* Works in both directions

We are not showing the rules for \Leftrightarrow , \oplus , and punctuation.

Rules for quantifiers and equality

We will introduce the following four rules.

▶ \forall -INTRO

▶ \forall -ELIM

▶ \exists -INTRO

▶ \exists -ELIM

We will also introduce rules for equality

▶ REFLEX

▶ EQSUB

Note

We will not show all steps due to propositional rules.

We will write 'propositional rules applied to ...'

Provably equivalent

Definition 15.1

*If statements $\{F\} \vdash G$ and $\{G\} \vdash F$ hold, then we say F and G are **provably equivalent**.*

Topic 15.2

Introduction rules for \forall and \exists

\exists -Intro quantifiers

If a fact is true about a term, we can introduce \exists

The condition is often not explicitly written. By writing $F(y)$ and $F(t)$, people may imply that the substitutions are defined.

$$\exists - \text{INTRO} \frac{\Sigma \vdash F(t)}{\Sigma \vdash \exists y. F(y)} \quad y \notin FV(F(z)), F(z)\{z \mapsto t\} \text{ and } F(z)\{z \mapsto y\} \text{ are defined}$$

for some variable z .

Example 15.1

1. $\{H(x)\} \vdash H(x)$
2. $\{H(x)\} \vdash \exists y. H(y)$

Assumption

\exists -Intro applied to 1

Bad derivations that violate the side condition $y \notin FV(F(z))$

Example 15.2

1. $\{1 \neq 2, x = 1, y = 2\} \vdash x \neq y$

Premise

2. $\{1 \neq 2, x = 1, y = 2\} \vdash \exists y. y \neq y$

\exists -Intro applied to 1 **X**

because $y \in FV(z \neq y)$.

Exercise 15.1

1. $\Sigma \vdash F(f(x), y)$

Premise

2. $\Sigma \vdash \exists y. F(y, y)$

\exists -Intro applied to 1 **X**

Give $F(z)$ that shows $y \in FV(F(z))$.

Bad derivation that violate the side condition ' $F(z)\{z \mapsto y\}$ is defined'

Example 15.3

1. $\{\exists y. c \neq y\} \vdash \exists y. c \neq y$

Assumption

2. $\{\exists y. c \neq y\} \vdash \exists y. \exists y. y \neq y$

\exists -Intro applied to 1X

because $(\exists y. z \neq y)\{z \mapsto y\}$ is not defined.

The following derivation is correct even if y is quantified somewhere in the formula.

Exercise 15.2

1. $\Sigma \vdash \exists w. (c \neq w \wedge \forall y. P(y))$

Assumption

2. $\Sigma \vdash \exists y. \exists w. (y \neq w \wedge \forall y. P(y))$

\exists -Intro applied to 1✓

Give $F(z)$ that shows all conditions are satisfied.

Commentary: In the first example, y being quantified is not solely responsible. The problem is that z is occurring in a scope where where y is quantified.

Bad derivations that violate the side condition ' $F(z)\{z \mapsto t\}$ is defined'

Example 15.4

1. $\Sigma \vdash \forall x. f(x) = x$

2. $\Sigma \vdash \exists y \forall x. y = x$

Statement 2 says that the domain is singleton, which is not implied by 1

Premise

\exists -Intro applied to 1

because $(\forall x. z = x)\{z \mapsto f(x)\}$ is not defined.

We get $F(t)$, we need to identify $F(z)$.

Commentary: z is a placeholder. $F(z)$ neither occurs in antecedents nor in consequent of the proof rule. Therefore, it is our choice (the person who is writing the proof) to choose z and $F(z)$. If we choose a z that is already around, then we may potentially run into a situation where some actions are not allowed. Therefore, it is cleaner to assume z is not being used for any other purpose in the context. Therefore, we should always choose such that z is not quantified in $F(z)$. If we choose $F(z)$ poorly, we may not be able to apply the rule.

Good derivations that may look bad

Not all occurrences of t are replaced.

One may complain that not all copies of $g(c)$ were replaced.

Example 15.5

1. $\emptyset \vdash \exists x_2. f(g(c), x_2) = f(g(c), c)$

2. $\emptyset \vdash \exists x_1, x_2. f(x_1, x_2) = f(g(c), c)$

$F(z) = \exists x_2. f(z, x_2) = f(g(c), c)$ satisfies all the side conditions.

Premise

\exists -Intro applied to 1 ✓

How to intro \forall ?

We have seen the following proof in our life.

- ▶ Consider a **fresh** name x to represent a number.
- ▶ We prove $Fact(x)$
- ▶ We conclude $\forall x.Fact(x)$.

\forall -Intro for variables

If something is true about a **variable** that is **not referred** elsewhere.

Then it must be true for any value in the universe.

No reference condition

$$\forall - \text{INTRO} \frac{\Sigma \vdash F(x)}{\Sigma \vdash \forall y. F(y)} \quad y \notin FV(F(z)), \quad x, z \in \text{Vars}, \quad \text{and } x \notin FV(\Sigma \cup \{F(z)\}).$$

Example 15.6

1. $\{H(x)\} \vdash H(x)$
2. $\{H(x)\} \vdash \forall y. H(y)$

Assumption

\forall -Intro applied to 1X

Since x is referred in left hand side, the above derivation is wrong.

Exercise 15.3

Why $FV(F(z))$ must not contain x ?

Commentary: The rule has implicit side condition that $F(z)\{z \mapsto x\}$ and $F(z)\{z \mapsto y\}$ are defined.

\forall -Intro (for constants)

Constants may play the similar role

$$\forall - \text{INTRO} \frac{\Sigma \vdash F(c)}{\Sigma \vdash \forall y. F(y)} \quad y \notin FV(F(z)), c \text{ is not referred in } \Sigma \cup \{F(z)\}, \text{ and } c/0 \in \mathbf{F},$$

for some variable z .

Example 15.7

1. $\Sigma \vdash H(c)$
2. $\Sigma \vdash \forall y. H(y)$

Premise and c is not referred in Σ

\forall -Intro applied to 1

Example: Bad \forall -Intro

Example 15.8

Consider the following derivation where we used a term for \forall -INTRO.

1. $\emptyset \vdash \exists y. f(y) \neq y \vee f(c) = c$
2. $\emptyset \vdash \forall x. (\exists y. f(y) \neq y \vee x = c)$

Premise

\forall -INTRO applied to 1 **X**

Our $F(z) = \exists y. f(y) \neq y \vee z = c$.

$f(c)$ does not occur in $F(z)$.

The formula in 1 is a valid formula and the formula in 2 is not a valid formula.

Topic 15.3

Elimination rules for \forall and \exists

Universal instantiation

If some thing is **always true**, we should be able to make **it true on any value**.

$$\forall - \text{ELIM} \frac{\Sigma \vdash \forall x. F(x)}{\Sigma \vdash F(t)}$$

Our first FOL proof : \forall implies \exists

Theorem 15.1

If we have $\Sigma \vdash \forall x.F(x)$, we can derive $\Sigma \vdash \exists x.F(x)$.

Proof.

1. $\Sigma \vdash \forall x.F(x)$
2. $\Sigma \vdash F(x)$
3. $\Sigma \vdash \exists x.F(x)$

the proof does not work
in the reverse direction

Premise

\forall -Elim applied to 1

\exists -Intro applied to 2

□

Exercise 15.4

Show $\Sigma \vdash \forall x.(F(x) \wedge G(x))$ and $\Sigma \vdash \forall x.F(x) \wedge \forall x.G(x)$ are provably equivalent.

One more example: working with quantifiers

Example 15.9

A derivation for $\emptyset \vdash (\forall x. (P(x) \vee Q(x)) \Rightarrow \exists x.P(x) \vee \forall x.Q(x))$.

1. $\{\forall x. (P(x) \vee Q(x)), \neg\exists x.P(x)\} \vdash \forall x. (P(x) \vee Q(x))$ *Assumption*
 2. $\{\forall x. (P(x) \vee Q(x)), \neg\exists x.P(x)\} \vdash P(y) \vee Q(y)$ *\forall -Elim applied to 1*
 3. $\{\forall x. (P(x) \vee Q(x)), \neg\exists x.P(x)\} \vdash \neg\exists x.P(x)$ *Assumption*
 4. $\{\forall x. (P(x) \vee Q(x)), \neg\exists x.P(x), P(y)\} \vdash P(y)$ *Assumption*
 5. $\{\forall x. (P(x) \vee Q(x)), \neg\exists x.P(x), P(y)\} \vdash \exists x.P(x)$ *\exists -Intro applied to 4*
 6. $\{\forall x. (P(x) \vee Q(x)), \neg\exists x.P(x)\} \vdash Q(y)$ *propositional rules applied to 2, 3, and 5*
 7. $\{\forall x. (P(x) \vee Q(x)), \neg\exists x.P(x)\} \vdash \forall x.Q(x)$ *\forall -Intro applied to 6*
- rest is propositional reasoning

Exercise 15.5

Fill the gaps in the step 6 and the tail of the proof.

Commentary: To understand the interplay of propositional reasoning and quantifiers, please solve the above exercise.

How to understand substitutions in the proof rules?

In the proof rules, there is a leaving term t and an arriving term s , and there is $F(z)$.
Antecedents have $F(t)$ and consequences have $F(s)$.

For example,

$$F(z) = \underbrace{P(z) \wedge \forall z.Q(z) \wedge (\forall w.R(w, u) \vee \exists y.R(z, y))}_{\text{No worry occurrences of } z}$$

Commentary: A good way to think is that the name of a quantified variable is not important to the outside world, except when we try to substitute a free variable in its scope by a term, which may have a variable with the same name.

The name conflict issue is a mute point. As long as we follow some naming discipline, which ensures that free variables in a system and quantified variables do not 'clash'. We need not worry. This is often done in programming languages. For example, import in python prefixes every imported name.

There are four cases of occurrences of z .

- ▶ z may occur free under no scope
- ▶ z is quantified in a scope
- ▶ free z does not occur in scope of a quantifier w
- ▶ free z occurs in scope of a quantifier y

(troubling case)

Only the last case causes a restriction that t and s cannot have y .

Where is \exists instantiation?

\exists can not behave like \forall .

If there is something, should we not be able to choose it? **Not an arbitrary** choice.

Example 15.10

Let us suppose we want to prove, "If there is a door in the building, I can steal diamonds."

Intuitively, we do...

1. Assume door x is there
2. \vdots
3. details of robbery
4. \vdots
5. I steal diamonds.
6. We say, therefore the theorem holds.

Formally, we need to do the following.

1. $\Sigma \cup \{D(x)\} \vdash D(x)$ *Assumption*
2. \vdots
3. symbolic details of robbery
4. \vdots
5. $\Sigma \cup \{D(x)\} \vdash Stolen$...
6. $\Sigma \vdash D(x) \Rightarrow Stolen$ \Rightarrow -Intro applied to 5
7. $\Sigma \vdash \exists x.D(x) \Rightarrow Stolen$ *What rule?*

Instantiation rule for exists

The following rule plays the role of \exists instantiation.

$$\exists - \text{ELIM} \frac{\Sigma \vdash F(x) \Rightarrow G}{\Sigma \vdash \exists y.F(y) \Rightarrow G} \quad x \notin FV(\Sigma \cup \{G, F(z)\}), y \notin FV(F(z))$$

Example: using \exists -Elim

Example 15.11

The following derivation proves $\emptyset \vdash \exists x.(A(x) \wedge B(x)) \Rightarrow \exists x.A(x)$

- | | |
|--|-----------------------------------|
| 1. $\{A(x) \wedge B(x)\} \vdash A(x) \wedge B(x)$ | Assumption |
| 2. $\{A(x) \wedge B(x)\} \vdash A(x)$ | \wedge -Elim applied to 1 |
| 3. $\{A(x) \wedge B(x)\} \vdash \exists x. A(x)$ | \exists -Intro applied to 2 |
| 4. $\emptyset \vdash A(x) \wedge B(x) \Rightarrow \exists x. A(x)$ | \Rightarrow -Intro applied to 3 |
| 5. $\emptyset \vdash \exists x.(A(x) \wedge B(x)) \Rightarrow \exists x. A(x)$ | \exists -Elim applied to 4 |

We cannot instantiate \exists out of the blue. We assume instantiated formula (step 1), prove the goal (step 3), and produce an implication (step 4), which is followed by \exists -Elim.

Exercise 15.6

Show $\Sigma \vdash \exists x.(F(x) \vee G(x))$, and $\Sigma \vdash \exists x.F(x) \vee \exists x.G(x)$ are provably equivalent.

Example: Disastrous derivations

Example 15.12

Here are two derivations that apply proof rules incorrectly and derive a bad statement.

1. $\{A(x)\} \vdash A(x)$ *Assumption*
2. $\{A(x)\} \vdash \forall x. A(x)$ *\forall -Intro applied to 1* **X**
3. $\emptyset \vdash A(x) \Rightarrow \forall x. A(x)$ *\Rightarrow -Intro applied to 2*
4. $\emptyset \vdash \exists x. A(x) \Rightarrow \forall x. A(x)$ *\exists -Elim applied to 3*

1. $\{\exists x. A(x)\} \vdash \exists x. A(x)$ *Assumption*
2. $\{\exists x. A(x)\} \vdash A(x)$ *\exists -Elim applied to 1* **X**
3. $\{\exists x. A(x)\} \vdash \forall x. A(x)$ *\forall -Intro applied to 2*
4. $\emptyset \vdash \exists x. A(x) \Rightarrow \forall x. A(x)$ *\Rightarrow -Intro applied to 3*

Topic 15.4

Rules for equality

Equality rules

For equality

$$\text{REFLEX} \frac{}{\Sigma \vdash t = t} \qquad \text{EQSUB} \frac{\Sigma \vdash F(t) \quad \Sigma \vdash t = t'}{\Sigma \vdash F(t')}$$

Exercise 15.7

Do we need a side condition for rule EQSUB?

Example : example for equality

Example 15.13

Let us prove $\emptyset \vdash \forall x, y. (x \neq y \vee f(x) = f(y))$

1. $\{x = y\} \vdash x = y$
2. $\{x = y\} \vdash f(x) = f(x)$
3. $\{x = y\} \vdash f(x) = f(y)$
4. $\{\} \vdash \neg(x = y) \vee f(x) = f(y)$
5. $\{\} \vdash \forall x, y. (\neg(x = y) \vee f(x) = f(y))$

Assumption

Reflex

EqSub applied to 1 and 2

propositional rules applied to 3

\forall -Intro applied twice to 4

Exercise 15.8

Write $F(z)$ s in the application of \forall -Intro.

Deriving symmetry for equality

Theorem 15.2

If we have $\Sigma \vdash s = t$, we can derive $\Sigma \vdash t = s$

Proof.

1. $\Sigma \vdash s = t$

Premise

2. $\Sigma \vdash s = s$

Reflex

3. $\Sigma \vdash t = s$

EqSub applied to 2 and 1 where $F(z) = (z = s)$

□

Therefore, we declare the following as a derived proof rule.

$$\text{EQSYMM} \frac{\Sigma \vdash s = t}{\Sigma \vdash t = s}$$

Example : finding evidence of \exists is hard

There are magic terms that can provide evidence of \exists . Here is an extreme example.

Example 15.14

Consider $\emptyset \vdash \exists x_4, x_3, x_2, x_1. f(x_1, x_3, x_2) = f(g(x_2), j(x_4), h(x_3, a))$

Let us construct a proof for the above as follows

1. $\emptyset \vdash f(g(h(j(c), a)), j(c), h(j(c), a)) = f(g(h(j(c), a)), j(c), h(j(c), a))$ *Reflex*
2. $\emptyset \vdash \exists x_1. f(x_1, j(c), h(j(c), a)) = f(g(h(j(c), a)), j(c), h(j(c), a))$ *\exists -Intro applied to 1*
3. $\emptyset \vdash \exists x_2. \exists x_1. f(x_1, j(c), x_2) = f(g(x_2), j(c), h(j(c), a))$ *\exists -Intro applied to 2*
4. $\emptyset \vdash \exists x_3. \exists x_2. \exists x_1. f(x_1, x_3, x_2) = f(g(x_2), j(c), h(x_3, a))$ *\exists -Intro applied to 3*
5. $\emptyset \vdash \exists x_4. \exists x_3. \exists x_2. \exists x_1. f(x_1, x_3, x_2) = f(g(x_2), j(x_4), h(x_3, a))$ *\exists -Intro applied to 4*

Topic 15.5

Problems

Exercise: extended \forall -elim rule

Exercise 15.9

Show that the following derived rule is sound

$$\forall - \text{ELIM} \frac{\Sigma \vdash \forall x_1 \dots x_n. F}{\Sigma \vdash F\sigma} \quad F \text{ is quantifier-free}$$

Exercise 15.10

Show that the following derived rule is sound

$$\forall - \text{SUBST} \frac{\Sigma \vdash \forall x_1 \dots x_n. F}{\Sigma \vdash \forall \text{Vars}(F\sigma). F\sigma} \quad F \text{ is quantifier-free and } FV(\Sigma) = \emptyset$$

Exercise : derived rules for equality

Exercise 15.11

Prove the following derived rules

$$\text{EQTRANS} \frac{\Sigma \vdash s = t \quad \Sigma \vdash t = r}{\Sigma \vdash s = r}$$

$$\text{PARAMODULATION} \frac{\Sigma \vdash s = t}{\Sigma \vdash r(s) = r(t)}$$

Practice formal proofs

Exercise 15.12

Prove the following statements

1. $\emptyset \vdash \forall x. \exists y. \forall z. \exists w. (R(x, y) \vee \neg R(w, z))$
2. $\emptyset \vdash \forall x. \exists y. x = y$
3. $\emptyset \vdash \forall x. \forall y. ((x = y \wedge f(y) = g(y)) \Rightarrow (h(f(x)) = h(g(y))))$
4. $\emptyset \vdash \exists x_1, x_2, x_3. f(g(x_1), x_2) = f(x_3, x_1)$

Proofs on set theory**

Exercise 15.13

Consider the following axioms of set theory

$$\Sigma = \{ \forall x, y, z. ((z \in x \Leftrightarrow z \in y) \Rightarrow x = y), \\ \forall x, y. (x \subseteq y \Leftrightarrow \forall z. (z \in x \Rightarrow z \in y)), \\ \forall x, y, z. (z \in x - y \Leftrightarrow (z \in x \wedge z \notin y)) \}.$$

Prove the following

$$\Sigma \vdash \forall x, y. x \subseteq y \Rightarrow \exists z. (y - z = x)$$

Exercise: bad orders

Exercise 15.14

Prove that the following formulas are mutually unsatisfiable.

- ▶ $\forall x. \neg E(x, x)$
- ▶ $\forall x, y. (E(x, y) \wedge E(y, x) \Rightarrow x = y)$
- ▶ $\forall x, y, z. (E(x, y) \wedge E(y, z) \Rightarrow \neg E(x, z))$
- ▶ $\forall x, y, z. (E(x, y) \wedge E(x, z) \Rightarrow E(y, x) \vee E(z, y))$
- ▶ $\exists x, y. E(x, y)$

Exercise: modeling equality using a predicate and axioms

Exercise 15.15

1. Give a formal proof that shows that following formulas are mutually unsatisfiable.

- ▶ $\forall x, y. x = y$
- ▶ $\forall x. \neg R(x, x)$
- ▶ $\exists x, y. R(x, y)$

2. Give a model that satisfies the following set of formulas.

- ▶ $\forall x. E(x, x)$
- ▶ $\forall x, y. E(x, y)$
- ▶ $\forall x, y. (E(x, y) \Rightarrow E(y, x))$
- ▶ $\forall x. \neg R(x, x)$
- ▶ $\forall x, y, z. (E(x, y) \wedge E(y, z) \Rightarrow E(x, z))$
- ▶ $\exists x, y. R(x, y)$

3. Give a formal proof that shows that the following formulas are mutually unsatisfiable.

- ▶ $\forall x. E(x, x)$
- ▶ $\forall x, y. E(x, y)$
- ▶ $\forall x, y. (E(x, y) \Rightarrow E(y, x))$
- ▶ $\forall x. \neg R(x, x)$
- ▶ $\forall x, y, z. (E(x, y) \wedge E(y, z) \Rightarrow E(x, z))$
- ▶ $\exists x, y. R(x, y)$
- ▶ $\forall x_1, x_2, y_1, y_2. (E(x_1, x_2) \wedge E(y_1, y_2) \wedge R(x_1, y_1) \Rightarrow R(x_2, y_2))$

Exercise: different proof systems (midterm 2021)

Exercise 15.16

Let us suppose we remove $\forall - \text{ELIM}$ from our FOL proof system and we add the following proof rule in our proof system.

$$\exists - \text{DEF} \frac{\Sigma \vdash \forall x.F(x)}{\Sigma \vdash \neg \exists x. \neg F(x)}$$

Show that we can drive $\forall - \text{ELIM}$ from the modified proof system. Give detailed derivation without skipping any step. Only formal derivations will be accepted.

Commentary: **Solution:**

- | | |
|---|-------------------------------|
| 1. $\Sigma \vdash \forall x.F(x)$ | Premise |
| 2. $\Sigma \cup \{\neg F(t)\} \vdash \forall x.F(x)$ | Monotonic applied to 1 |
| 3. $\Sigma \cup \{\neg F(t)\} \vdash \neg \exists x. \neg F(x)$ | \exists -Def applied to 1 |
| 4. $\Sigma \cup \{\neg F(t)\} \vdash \neg F(t)$ | Assumption |
| 5. $\Sigma \cup \{\neg F(t)\} \vdash \exists x. \neg F(x)$ | \exists -Intro applied to 4 |
| 6. $\Sigma \vdash \neg \neg F(t)$ | ByContra applied to 3 and 5 |
| 7. $\Sigma \vdash F(t)$ | RevDoubleNeg applied to 6 |

Proofs on arrays (midterm 2022)

Exercise 15.17

Let Σ contain the following FOL sentences (all free symbols are functions or constants)

1. $\forall z, i, x. \text{read}(\text{store}(z, i, x), i) = x$
2. $\forall z, i, j, v. (i = j \vee \text{read}(\text{store}(z, i, v), j) = \text{read}(z, j))$
3. $\text{store}(a, n, \text{read}(b, n)) = \text{store}(b, n, \text{read}(a, n))$
4. $\text{read}(b, m) \neq \text{read}(a, m)$

Using the formal proof system, show that Σ can derive contradiction.

Commentary: Solution: The following proof is repetitive. Key observation is what to substitute for v and x and aim to derive $m = n$.

1. $\Sigma \vdash \text{store}(a, n, \text{read}(b, n)) = \text{store}(b, n, \text{read}(a, n))$ Assumption
2. $\Sigma \vdash \forall z, i, j, v. (i = j \vee \text{read}(\text{store}(z, i, v), j) = \text{read}(z, j))$ Assumption
3. $\Sigma \vdash (n = m \vee \text{read}(\text{store}(a, n, \text{read}(b, n)), m) = \text{read}(a, m))$ \forall -Elim applied to 1 with substitutions $\{z \mapsto a, i \mapsto n, j \mapsto m, v \mapsto \text{read}(b, n)\}$
4. $\Sigma \vdash (n = m \vee \text{read}(\text{store}(b, n, \text{read}(a, n)), m) = \text{read}(b, m))$ \forall -Elim applied to 1 with substitutions $\{z \mapsto b, i \mapsto n, j \mapsto m, v \mapsto \text{read}(a, n)\}$
5. $\Sigma \vdash (n = m \vee \text{read}(\text{store}(b, n, \text{read}(a, n)), m) = \text{read}(a, m))$ EqSub applied to 3 and 1
6. $\Sigma \vdash (n = m \vee \text{read}(b, m) = \text{read}(a, m))$ EqSub applied to 3 and 5, and some propositional reasoning
7. $\Sigma \vdash \text{read}(b, m) \neq \text{read}(a, m)$ Assumption
8. $\Sigma \vdash n = m$ Resolution applied to 6 and 7
9. $\Sigma \vdash \forall z, i, x. \text{read}(\text{store}(z, i, x), i) = x$ Assumption
10. $\Sigma \vdash \text{read}(\text{store}(a, n, \text{read}(b, n)), n) = \text{read}(b, n)$ \forall -Elim applied to 9 with substitutions $\{z \mapsto a, i \mapsto n, x \mapsto \text{read}(b, n)\}$
11. $\Sigma \vdash \text{read}(\text{store}(b, n, \text{read}(a, n)), n) = \text{read}(a, n)$ \forall -Elim applied to 9 with substitutions $\{z \mapsto b, i \mapsto n, x \mapsto \text{read}(a, n)\}$
12. $\Sigma \vdash \text{read}(\text{store}(b, n, \text{read}(a, n)), n) = \text{read}(b, n)$ Eqsub applied to 10 and 11
13. $\Sigma \vdash \text{read}(b, n) = \text{read}(a, n)$ Eqsub applied to 11 and 12
14. $\Sigma \vdash \text{read}(b, m) = \text{read}(a, m)$ Eqsub applied to 13 and 8

Topic 15.6

Extra slides: Soundness

Soundness of the proof system

We need to show that the proof rules derive only valid statements.

We only need to prove the soundness of the new proof rules in addition to the propositional rule.

Substitution

Theorem 15.3

For a variable z , a term t , and a formula $F(z)$, if $m^\nu(z) = m^\nu(t)$ and $F(t)$ is defined, then

$$m, \nu \models F(z) \quad \text{iff} \quad m, \nu \models F(t).$$

Proof.

Not so trivial proof by structural induction. □

Exercise 15.18

Write down the above proof. Hint: You need to case split when we quantify over z or some other variable.

Soundness: \exists – INTRO is sound

Theorem 15.4

The following rule is sound.

$$\exists - \text{INTRO} \frac{\Sigma \vdash F(t)}{\Sigma \vdash \exists y. F(y)} \quad y \notin FV(F(z)), F(z)\{z \mapsto t\} \text{ and } F(z)\{z \mapsto y\} \text{ are defined}$$

for some variable z .

Proof.

1. Let us assume $m, \nu \models \Sigma$.
2. Due to the antecedent, $m, \nu \models F(t)$. Let $m^\nu(t) = v$.
3. Since $z \notin FV(F(t))$, $m, \nu[z \mapsto v] \models F(t)$.
4. Since $F(z)\{z \mapsto t\}$ is defined, $m, \nu[z \mapsto v] \models F(z)$. (why?) (Theorem 15.3)
5. Since $y \notin FV(F(z))$, $m, \nu[z \mapsto v, y \mapsto v] \models F(z)$.
6. Since $F(z)\{z \mapsto y\}$ is defined, $m, \nu[z \mapsto v, y \mapsto v] \models F(y)$. (Theorem 15.3)
7. Therefore, $m, \nu[z \mapsto v] \models \exists y. F(y)$.
8. Since $z \notin FV(F(y))$, $m, \nu \models \exists y. F(y)$ □

Commentary: All soundness proofs are repeated applications of similar arguments. However, in each rule the side conditions play their role differently. To understand the side conditions, please look into all the soundness arguments in the extra slides of this lecture.

Soundness: \forall – INTRO is sound

Theorem 15.5

The following rule is sound.

$$\forall - \text{INTRO} \frac{\Sigma \vdash F(x)}{\Sigma \vdash \forall y. F(y)} \quad y \notin FV(F(z)), \quad x, z \in \text{Vars}, \quad \text{and } x \notin FV(\Sigma \cup \{F(z)\}).$$

Proof.

- ▶ Let us assume $m, \nu \models \Sigma$. Let v be some value in the domain of model m .
- ▶ Since $x \notin FV(\Sigma)$, $m, \nu[x \mapsto v] \models \Sigma$. Due to the antecedent, $m, \nu[x \mapsto v] \models F(x)$.
- ▶ Since $z \notin FV(F(x))$, $m, \nu[x \mapsto v, z \mapsto v] \models F(x)$.
- ▶ Since $F(z)\{z \mapsto x\}$ is defined, $m, \nu[x \mapsto v, z \mapsto v] \models F(z)$ _(why?).
- ▶ Since $x \notin FV(F(z))$, $m, \nu[z \mapsto v] \models F(z)$.
- ▶ Since $y \notin FV(F(z))$, $m, \nu[z \mapsto v, y \mapsto v] \models F(z)$.
- ▶ Since $F(z)\{z \mapsto y\}$ is defined, $m, \nu[x \mapsto v, z \mapsto v] \models F(y)$ _(why?).
- ▶ Since $z \notin FV(F(y))$ _(why?), $m, \nu[y \mapsto v] \models F(y)$.
- ▶ Since v is an arbitrary value, we have $m, \nu \models \forall y. F(y)$.

□

Soundness: \forall – ELIM is sound

Theorem 15.6

The following rule is sound.

$$\forall - \text{ELIM} \frac{\Sigma \vdash \forall x. F(x)}{\Sigma \vdash F(t)}$$

Proof.

1. Let $t' = t\{x \mapsto z\}$, where z is a fresh variable.
2. Since $F\{x \mapsto t\}$ is defined, $F\{x \mapsto t'\}$ is defined and $F(t')\{z \mapsto x\}$ is defined.
3. Let us assume $m, \nu \models \Sigma$. Let $\nu' \triangleq \nu[z \mapsto \nu(x)]$. Since $z \notin FV(\Sigma)$, $m, \nu' \models \Sigma$.
4. Due to the antecedent, $m, \nu' \models \forall x. F(x)$.
5. Let $v \triangleq m^{\nu'}(t')$. Since $x \notin \text{Vars}(t')$, $v = m^{\nu'[x \mapsto v]}(t')$.
6. Due to \forall semantics, $m, \nu'[x \mapsto v] \models F(x)$.
7. Since $F\{x \mapsto t'\}$ is defined, $m, \nu'[x \mapsto v] \models F(t')$.
8. Since $x \notin FV(F(t'))$, $m, \nu' \models F(t')$.
9. Therefore, $m, \nu \models F(t)$._(why?)

Commentary: If x does not occur in t , the proof is simpler. However, it occurs very often in practice.

□

Soundness: \exists – ELIM is sound

Theorem 15.7

The following rule is sound.

$$\exists - \text{ELIM} \frac{\Sigma \vdash F(x) \Rightarrow G}{\Sigma \vdash \exists y.F(y) \Rightarrow G} \quad x \notin FV(\Sigma \cup \{G, F(z)\}), y \notin FV(F(z))$$

Proof.

- ▶ Let us assume $m, \nu \models \Sigma$ and $m, \nu \models \exists y.F(y)$.
- ▶ There is v in domain of m such that $m, \nu[y \mapsto v] \models F(y)$.
- ▶ Since $x, y \notin FV(F(z))$, and $F(x)$ and $F(y)$ substitutions are defined, $m, \nu[x \mapsto v] \models F(x)$.
- ▶ Since $x \notin FV(\Sigma)$, $m, \nu[x \mapsto v] \models \Sigma$.
- ▶ Due to the antecedent, $m, \nu[x \mapsto v] \models F(x) \Rightarrow G$.
- ▶ Therefore, $m, \nu[x \mapsto v] \models G$.
- ▶ Since $x \notin FV(G)$, $m, \nu \models G$.

□

End of Lecture 15