Randomized Computation

NSPACE = PSPACE = coPSPACE

oracle for: PH — Power (computation) but no proof of correctness (randomness) — "Most computing you do is pointless"

Coin: NP — Unintuitive, non-constructive proofs

One sided: NP = CoNP — Correct, but explicit witness of small size

P — Efficient computation

P/poly — L — space resource

One string was missing, magically owns more advice & captures all of coNP then ?

Randomized Computation

Eg: Find a quadratic non-residue

This is a functional problem

$\mathbb{Z}_p = \mathbb{Z} \bmod \text{prime } p$ (odd) — collection of remainders/ residue

- is a field — addition, multiplication, division (of course)
- ring (odd^n, mod^n)

Let $\exists\, a \in \mathbb{Z}_p^*$ is a quadratic residue if $\exists\, b \in \mathbb{Z}_p^*$ s.t. $b^2 = a$

Consider the following map:

$\Psi : \mathbb{Z}_p^* \longrightarrow \mathbb{Z}_p^*$

$x \longrightarrow x^2$

this is a homomorphism

$\Psi(x) = \Psi(-x)$



2 to 1 map

$\Rightarrow \exists\ \frac{p-1}{2}$ quadratic residues

Everything not in the image(Ψ) is a quadratic non residue (QNR).

$\frac{p-1}{2}$ qnrs.

So, if $\mathbb{Z}_p^*$ $\frac{1}{2}$ qnr $\frac{1}{2}$ qnr

& Given p, find a qnr in $\mathbb{F}_p^*$ or $\mathbb{Z}_p^*$ in $\text{poly}(\log p)$ time.

- Randomized algo — pick a random element
- Det algo — (Open)

first qnr: $O(\log^2 n)$ assuming GRH (not unconditional)

(size in bits)

Can you think of a way to answer density$_{\mathbb{Z}_p^*}$ to get a non-residue in $\text{poly}(\log p)$ steps?

Notation: Legendre Symbol

$\left(\frac{a}{p}\right) = \begin{cases} 0 & \text{if } a \equiv_p 0 \\ 1 & \text{if } a \text{ is qr} \\ -1 & \text{if } a \text{ is qnr} \end{cases}$

$\left(\frac{a}{p}\right) \equiv_p a^{\frac{p-1}{2}}$ when p is odd prime.

Note $a^{\frac{p-1}{2}} \equiv_p 1 \quad \forall\, a \in \mathbb{Z}_p^*$ gives 1 or -1.

What about decision problem?

① Primality Testing

$\text{PRIMES} = \{\langle n \rangle_x \mid n \text{ is prime}\}$

& Given n, check if its a prime? in binary

$\mathbb{Z}_n$ = residues/remainders modulo n. This is a ring not a field. $2 \times 3 = 0$ — we invert to $\mathbb{Z}_n^*$

$a^{n-1} \equiv_n 1 \quad \forall\, a \in \mathbb{Z}_n \setminus \{0\}$ iff n is a prime.

Extending quadratic residues to non-primes

$\Psi : \mathbb{Z}_n^* \to \mathbb{Z}_n$

$x \longmapsto x^2$

$a \in \text{Im}(\Psi) \leq \mathbb{Z}_n^*$ is a quad-residue ($\exists$)

$\exists\ b$ s.t. $a \equiv b^2 \bmod n$

Jacobi Symbol $n = p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$

$\left(\frac{a}{n}\right) \triangleq \prod_{i=1}^{m} \left(\frac{a}{p_i}\right)^{k_i}$

$a^{\phi(n)} \equiv_n 1 \quad a \in \mathbb{Z}_n^*$

Computing $\phi(n)$ ≡ factoring n (Executive) ↓ polynomial reducible

Properties of Jacobi Symbol

1. $a \equiv_n b \implies \left(\frac{a}{n}\right) = \left(\frac{b}{n}\right)$
2. $\left(\frac{ab}{n}\right) = \left(\frac{a}{n}\right)\left(\frac{b}{n}\right)$ — multiplicative
3. $\left(\frac{-1}{n}\right) = (-1)^{\frac{n-1}{2}}$ & What if $a \notin \mathbb{Z}_n^*$, $\left(\frac{a}{n}\right)$?
4. Quadratic reciprocity law: m, n odd pos. $\mathbb{Z}$. $\left(\frac{m}{n}\right) = (-1)^{\frac{m-1}{2}\cdot\frac{n-1}{2}} \left(\frac{n}{m}\right)$ Proved
5. $\left(\frac{2}{n}\right) = (-1)^{\frac{n^2-1}{8}}$ n - odd

Compute $\left(\frac{1001}{3333}\right)$

Conclusion: Jacobi symbol can be computed in $O(\log n \log^2 n)$ time

Randomized algo for PRIMALITY

n - odd composite $\exists\ a$ s.t. $\left(\frac{a}{n}\right) \neq_n a^{\frac{n-1}{2}}$

$S_n = \left\{ a \in \mathbb{Z}_n^* \;\middle|\; \left(\frac{a}{n}\right) = a^{\frac{n-1}{2}} \bmod n \right\}$ is a subgroup of $\mathbb{Z}_n^*$ (easy to verify)

$|S_n| \leq \frac{|\mathbb{Z}_n^*|}{2}$

Solovay-Strassen Algorithm

1. Pick $a \in \mathbb{Z}_n \setminus \{0\}$ randomly
2. Compute $\gcd(a, n)$
3. If $\text{GCD}(a,n) \neq 1$: output composite
4. If $\left(\frac{a}{n}\right) \neq a^{\frac{n-1}{2}} \bmod n$: output composite; else prime

If n is prime

$\Pr_a[\text{SS}(n) = \text{prime}] = 1$

n - composite

$\Pr_a[\text{SS}(n) = \text{Prime}] = \frac{|S_n|}{|\mathbb{Z}_n^*|} \leq \frac{1}{2}$

Work for all n not some/most n.

Polynomial Identity Testing

Q Given a multivariate poly $f \in \mathbb{F}[x_1, \ldots, x_n]$ test if $f \equiv 0$?

Polynomial-Identity $= \{ f \in \mathbb{F}[x_1, \ldots, x_n] \mid f \equiv 0 \}$

Representation of polynomials
- Arithmetic/Algebraic Circuits/ Determinants
- Such representations

JEE — Put value trick

- Size of the input
- Univariate instance.

- Univariate degree d polynomial has always d roots

$f = f_d x^d + f_{d-1} x^{d-1} + \cdots + f_0$

Multivariate extension of this

Schwartz Zippel / PIT Lemma — total-degree d.

$\Pr_{a \in S^n}[P(a) = 0] \leq \frac{d}{|S|}$

$(a_1, a_2, \ldots, a_n) \in S \times S \cdots S$

$\Pr_{r_1}[f_1 = 0] \leq \Pr[f_1 = 0 \mid r_1 \neq 0] + \Pr_{r_1}[r_1 = 0]$

$\leq \frac{d-1}{|S|} + \frac{1}{|S|} = \frac{d}{|S|}$

Combinatorial nullstellensatz

Matching

Issue with Fermat Primality Test

Note $N = p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$

$$\mathbb{Z}_N^* \cong \mathbb{Z}_{p_1^{k_1}}^* \otimes \mathbb{Z}_{p_2^{k_2}}^* \otimes \cdots \otimes \mathbb{Z}_{p_m^{k_m}}^*$$

CRT

$\mathbb{Z}_n^*$ is not-necc. cyclic group

In $\mathbb{Z}_p^*$ there exist a generator

$\alpha \in \mathbb{Z}_p^*$ st $\text{Ord}(\alpha) = p-1$

$\alpha^{p-1} = 1$ (and not smaller)

$n = pq$

$$\mathbb{Z}_n^* \cong \mathbb{Z}_p^* \otimes \mathbb{Z}_q^*$$

the max order of $\alpha \in \mathbb{Z}_n^*$

is $\text{lcm}(p-1, q-1)$

and not

Carmichael numbers

$n = 561$

$= 3 \cdot 11 \cdot 17$

$$\mathbb{Z}_{561}^* \cong \mathbb{Z}_3^* \otimes \mathbb{Z}_{11}^* \otimes \mathbb{Z}_{17}^*$$

max order here $\text{lcm}(2, 10, 16) = 80$.

and $80 \mid 560$, thus $\forall \alpha \in \mathbb{Z}_{561}^*$

satisfy $\alpha^{560} = 1 \mod 561$.

[Fact] There are ∞ many carmichael numbers.

$$\frac{|\mathbb{Z}_n^*|}{|\mathbb{Z}_n|} \to 1 \quad \text{as} \quad n \to \infty$$

Randomized Complexity Classes.

PTM: TM M with two trans^n f^n δ_0, δ_1

- Every step choosen ind. at random δ_0 or δ_1


(only accpt/Rej)

- $M(x)$ is a r.v. "output".
- $T: \mathbb{N} \to \mathbb{N}$, M stops in $T(|x|)$ steps, for any x, for any random choice.

We say PTM M decides $L \subseteq \{0,1\}^*$ in time $T(n)$ if $\forall x \in \{0,1\}^*$

① $\Pr[M(x) = L(x)] \geq 2/3$

② M halts in $T(|x|)$ regardless of random choice.



BPP

$BPTIME(T(n)) \equiv$ Lang. decidable by PTM in $O(T(n))$ time.

$$BPP = \bigcup_{c \geq 1} BPTIME(n^c).$$

Note: Consistency check we want things to work on all inputs.

Although any such variation for most inputs w/ or w/o randomness makes sense $\equiv$ average case complexity.

If Not a fan of NTM, here is another alternate (equiv) def^n of BPP.

> $L \in BPP$ if $\exists$ poly time (deterministic) TM M and $p: \mathbb{N} \to \mathbb{N}$ (poly-bounded) s.t. $\forall x \in \{0,1\}^*$,
>
> $$\Pr_{r \in_R \{0,1\}^{p(|x|)}}[M(x,r) = L(x)] \geq 2/3$$

Q// What the error in yes/no case for PIT / Primality?

In SS, if n is a prime (yes instance) No error

In PIT, s

This motivates prob. algorithms / classes. w/ one sided error.

RTIME(T(n)): $L \in RTIME(T(n))$ if there is a PTM M running in time $T(n)$ s.t.

$$x \in L \implies \Pr(M(x) = 1) \geq 2/3$$

$$x \notin L \implies \Pr[M(x) = 1] = 0$$

$$RP = \bigcup_{c > 0} RTIME(n^c).$$

$$coRP = \{L \mid \bar{L} \in RP\}$$

Equiv: Error free on accpt, some error on reject.

Zero-sided Error


For a PTM M,

Let $T_{M,x}$ be a r.v.

$\Pr_{\text{random choice}}[T_{M,x} = T] = p$ if w/ prob p over randomness, M will halt on x in $T(|x|)$ steps.

▷ If no error and Expected run time $O(T(|x|)) \equiv ZTIME(T(n))$

$$ZPP = \bigcup_{c > 1} ZTIME(n^c).$$



In fact, Theorem $ZPP = RP \cap co\text{-}RP$. (Exercise)

Error Reduction in BPP.

$$BPP_{\frac{1}{2}+n^{-c}} = BPP_{2/3} = BPP_{\frac{1}{2}+2^{-n}}$$

≡ with succ. prob.

We can transform a machine w.s.p. $\frac{1}{2}+n^{-c}$ into machine w.s.p $2/3$. We can convert it to exponentially close to 1.

Thm: Let $L \subseteq \{0,1\}^*$ decidable by PTM M (polytime)

s.t. $\forall\, x \in \{0,1\}^*$

$$Pr\left[M(x) = L(x)\right] \geq \frac{1}{2} + |x|^{-c}$$

Then for every $d > 0$ (const.) $\exists$ PTM M'

s.t. $\forall\, x \in \{0,1\}^*$, $Pr\left[M'(x) = L(x)\right] \geq 1 - 2^{-|x|^d}$

M' does the following

Proof: Run $M(x)$ for $k = 8|x|^{2c+d}$ times and take Majority.

say $y_1, y_2, \ldots, y_k \in \{0,1\}$

$$X_i = \begin{cases} 1 & \text{if } y_i = L(x) \\ 0 & \text{ow.} \end{cases}$$

$$E[X_i] = Pr[X_i = 1] \geq p \quad \left(p = \frac{1}{2} + |x|^{-c}\right)$$

We want to analyze prob: $Pr_{y_1,\ldots,y_k}\left[\text{Majority}(y_1,\ldots,y_k) \neq L(x)\right]$

$\Updownarrow$

$\sum X_i < \frac{k}{2}$

$\sum X_i - \left(\frac{1}{2} + \frac{1}{n^{10}}\right)k$

$k\Delta > \frac{k}{n^{10}}$

Chernoff Bounds

$X_1, \ldots, X_n$ be ind. r.v. over $\{0,1\}$

Let $\mu = \sum_{i=1}^{n} E[X_i]$ and let $\mu = \sum_{i=1}^{n} E[X_i]$

Then for every/any $c > 0$

$$Pr\left[\left|\sum_{i=1}^{n} X_i - \mu\right| \geq c\mu\right] \leq 2e^{-\frac{c^2}{4}\mu}$$

set $\delta = |x|^{-c/2}$ guarantees that,


$$\sum_{i=1}^{k} X_i \geq pk - \delta pk$$

with err prob bounded by

$$e^{-\frac{1}{4|x|^2}\cdot\frac{1}{2}\cdot 8|x|^{2c+d}} \leq 2^{-|x|^d}. \quad \square$$

That is Pr of deviating from mean is $2^{-\Omega(\mu)}$

Bell curve



Q// Does the universe has endless stream of randomness?

↓

perfect?

$BPP \subseteq P_{/poly}$

- All BPP lang. have poly sized ckt.
- Big Picture: unless PH collapses not randomized poly-time algo for SAT.

Theorem 7.14 $BPP \subseteq P_{/poly}$

Proof: $L \in BPP$

$\Updownarrow$

$\exists$ TM M on inputs of size n, uses $m = poly(n)$ random bits s.t.

$$Pr_{r \in \{0,1\}^m}[M(x,r) = L(x)] \geq 2/3$$

(By error reduction via Majority) $1 - \frac{1}{2^{n+1}}$

Bad string $r \in \{0,1\}^m$

$M(x,r) \neq L(x)$.

Good string $r \in \{0,1\}^m$

$M(x,r) \equiv L(x)$.

bad strings for x (fixed) $\leq \frac{2^m}{2^{n+1}}$.

bad strings for any x. $\leq 2^n \cdot \frac{2^m}{2^{n+1}} \leq \frac{2^m}{2}$.

$\Rightarrow$ $\exists$ (say r_0) $r \in \{0,1\}^m$ that is good for every input of length n.

Cook Levin $\Rightarrow$ $M(x, r_0)$ is the required poly-sized ckt.

$BPP \subseteq \Sigma_2 \cap \Pi_2$

Set-size Estimation algorithm
(in $\Sigma_2 \cap \Pi_2$)

$L \in BPP$

$\Downarrow$ Error Red^n

$x \in L \implies \Pr_r [M(x,r) = 1] \geq 1 - 2^{-n}$

$x \notin L \implies \Pr_r [M(x,r) = 1] \leq 2^{-n}$

For any (fixed) $x \in \{0,1\}^n$,

$$S_x = \{ r \in \{0,1\}^m \mid M(x,r) = 1 \}$$

Then either

$|S_x| \geq (1-2^{-n}) 2^m$ or $|S_x| \leq 2^{-n} 2^m$.

For $S \subseteq \{0,1\}^m$ and $u \in \{0,1\}^m$

$S + u := \{ x + u : x \in S \}$

$+ = \oplus$ mod 2.

$k = \lceil \frac{m}{n} \rceil + 1$

Claim 1: If $|S| \leq 2^{m-n}$

k vectors $u_1, \ldots, u_k$ (any)

$$\bigcup_{i=1}^{k} (S + u_i) \neq \{0,1\}^m$$

Claim 2: For every set $S \subseteq \{0,1\}^m$ with $|S| \geq (1-2^{-n})2^m$, there exists $u_1, \ldots, u_k$ s.t. $\bigcup_{i=1}^{k} (S+u_i) = \{0,1\}^m$.

Proof: $u_1, \ldots, u_k \rightsquigarrow$ uni random $\in \{0,1\}^m$

$$Pr\left[\bigcup_{i=1}^{k} (S+u_i) = \{0,1\}^m\right] > 0$$

$B_r \equiv$ Bad even that $r \in \{0,1\}^m$ is not covered by $S+u_i$'s

$$r \notin \bigcup_{i=1}^{k} (S+u_i).$$

T.S.T $Pr\left[\exists_{r \in \{0,1\}^m} B_r\right] < 1$

$\Uparrow$ union bound

Note $\forall r \quad Pr[B_r] < 2^{-m}$.

$$B_r = \bigcap_{i \in [k]} B_r^i$$

$B_r^i \equiv r \notin S+u_i \iff r+u_i \notin S$ over $\mathbb{F}_2$

But $r+u_i$ is a random element of $\{0,1\}^m$

$$\Rightarrow Pr[B_r^i] \leq 2^{-n}$$

$$\Rightarrow Pr[B_r] = \left(Pr[B_r^i]\right)^k \leq 2^{-nk} < 2^{-m}.$$

Thus Σ_2 machine: $x \in L \iff$

$$\exists u_1, u_2 \cdots u_k \in \{0,1\}^m \ \forall r \in \{0,1\}^m$$

$$r \in \bigcup_{i=1}^{k} (S_x + u_i)$$

$$\Updownarrow$$

$$\exists u_1, \ldots u_k \in \{0,1\}^m \ \forall r \in \{0,1\}^m$$

$$\bigvee_{i=1}^{k} M(x, r \oplus u_i) \text{ accepts.}$$

Complete problems for BPP ?

BPTIME hierarchy ?


Semantic defn / Syntatic defn

Randomized Reductions

Def^n $B \leq_R C$ (Transitive?)

iff PTM M s.t. poly-time M.

$x \in \{0,1\}^*$, $\Pr[B(M(x)) = C(x)] \geq 2/3$.



$x \longmapsto M(x)$

Def^n: $BP \cdot NP = \{L : L \leq_R 3SAT\}$

Ex: $\overline{3SAT} \in BP \cdot NP$?

Randomized space-bounded Comp^n

BPL

$O(\log n)$-spaced PTM M s.t.

$$\Pr[M(x) = L(x)] \geq 2/3.$$

RL

$O(\log n)$-spaced PTM M s.t.

$$x \in L \Rightarrow \Pr[M(x) = 1] \geq 2/3$$

$$x \notin L \Rightarrow \Pr[M(x) = 1] = 0.$$

Thm: UPATH $\in$ RL.

Note, $RL \subseteq NL \subseteq P$

Is $BPL \subseteq P$? Ex

UPATH $\in$ RL: (Ex?)

Take a random walk of length $\ell = 100\, n^{10}$ from s.

Note it is known that,

$$BPL \subseteq L^{3/2} \quad \text{(Saks-Zhou)}$$

Project.